



8 The time-dependent Schrödinger equation

Slides: Lecture 8a Introduction to the time-dependent Schrödinger equation

Text reference: Quantum Mechanics for Scientists and Engineers

Chapter 3 introduction






The time-dependent Schrödinger equation

Quantum mechanics for scientists and engineers

David Miller





8 The time-dependent Schrödinger equation

Slides: Lecture 8b Rationalizing the time-dependent Schrödinger equation

Text reference: Quantum Mechanics for Scientists and Engineers

Sections 3.1 – 3.2





The time-dependent Schrödinger equation



Rationalizing the time-dependent Schrödinger equation

Quantum mechanics for scientists and engineers

David Miller

Relation between energy and frequency

The relation between

energy

and

frequency

for

photons

is

$$E = h\nu = \hbar\omega$$

Relation between energy and frequency

The relation between

energy

and

frequency

for

quantum mechanics

is

$$E = h\nu = \hbar\omega$$

Rationalizing the time-dependent equation

We want a time-dependent wave equation

for a particle with mass m

with this relation $E = h\nu = \hbar\omega$ between energy and frequency

We might also reasonably want it to have plane wave solutions

e.g., of the form $\exp[i(kz - \omega t)]$

when we have some specific energy E

and when we are in a uniform potential

Rationalizing the time-dependent equation

Schrödinger postulated the time-dependent equation

$$-\frac{\hbar^2}{2m}\nabla^2\Psi(\mathbf{r},t)+V(\mathbf{r},t)\Psi(\mathbf{r},t)=i\hbar\frac{\partial\Psi(\mathbf{r},t)}{\partial t}$$

Note that for a uniform potential

e.g., $V = 0$ for simplicity

with $E = \hbar\omega$ and $k = \sqrt{2mE / \hbar^2}$

waves of the form

$$\exp\left[-i(\omega t \pm kz)\right] \equiv \exp\left[-i\left(\frac{Et}{\hbar} \pm kz\right)\right] \equiv \exp\left(-i\frac{Et}{\hbar}\right)\exp(\mp ikz)$$

are indeed solutions

Rationalizing the time-dependent equation

In his time-dependent equation

$$-\frac{\hbar^2}{2m}\nabla^2\Psi(\mathbf{r},t)+V(\mathbf{r},t)\Psi(\mathbf{r},t)=i\hbar\frac{\partial\Psi(\mathbf{r},t)}{\partial t}$$

Schrödinger chose a sign for the right hand side
which means that a wave with a spatial part

$$\propto \exp(ikz)$$

is definitely going in the positive z direction

That wave, including its time dependence
would be of the form (for $V = 0$)

$$\exp\left[i(kz - Et / \hbar)\right]$$

Compatibility with the time-independent equation

Before examining the time-dependent equation further
first we should check that it is compatible with the
time-independent equation

The time-independent equation could apply if we had
states of definite energy E , an eigenenergy

Suppose we had some corresponding eigenfunction $\psi(\mathbf{r})$
so that

$$-\frac{\hbar^2}{2m}\nabla^2\psi(\mathbf{r}) + V(\mathbf{r})\psi(\mathbf{r}) = E\psi(\mathbf{r})$$

Compatibility with the time-independent equation

As it stands

this solution $\psi(\mathbf{r})$ is not a solution of the time-dependent equation

$$-\frac{\hbar^2}{2m}\nabla^2\Psi(\mathbf{r},t)+V(\mathbf{r},t)\Psi(\mathbf{r},t)=i\hbar\frac{\partial\Psi(\mathbf{r},t)}{\partial t}$$

Putting $\psi(\mathbf{r})$ in here for $\Psi(\mathbf{r},t)$ does not work

because $\psi(\mathbf{r})$ has no time-dependence

the right hand side is zero

whereas it should be $E\psi(\mathbf{r})$

how do we resolve this?

Compatibility with the time-independent equation

Suppose that, instead of proposing the solution $\psi(\mathbf{r})$

we propose $\Psi(\mathbf{r}, t) = \psi(\mathbf{r}) \exp(-iEt / \hbar)$

$$-\frac{\hbar^2}{2m} \nabla^2 \Psi(\mathbf{r}, t) + V(\mathbf{r}) \Psi(\mathbf{r}, t)$$

$$= -\frac{\hbar^2}{2m} \nabla^2 \psi(\mathbf{r}) \exp(-iEt / \hbar) + V(\mathbf{r}) \psi(\mathbf{r}) \exp(-iEt / \hbar)$$

$$= \left[-\frac{\hbar^2}{2m} \nabla^2 \psi(\mathbf{r}) + V(\mathbf{r}) \psi(\mathbf{r}) \right] \exp(-iEt / \hbar) = E \psi(\mathbf{r}) \exp(-iEt / \hbar)$$

$$= E \Psi(\mathbf{r}, t)$$

so $\Psi(\mathbf{r}, t) = \psi(\mathbf{r}) \exp(-iEt / \hbar)$ **solves the time-independent Schrödinger equation**

Compatibility with the time-independent equation

Similarly, knowing that $\psi(\mathbf{r})$ solves the time-independent equation with energy E

substituting $\Psi(\mathbf{r}, t) = \psi(\mathbf{r}) \exp(-iEt / \hbar)$
in the time-dependent equation gives

$$\begin{aligned} -\frac{\hbar^2}{2m} \nabla^2 \Psi(\mathbf{r}, t) + V(\mathbf{r}) \Psi(\mathbf{r}, t) &= i\hbar \frac{\partial \Psi(\mathbf{r}, t)}{\partial t} = i\hbar \frac{\partial}{\partial t} [\psi(\mathbf{r}) \exp(-iEt / \hbar)] \\ &= i\hbar \psi(\mathbf{r}) \frac{\partial}{\partial t} [\exp(-iEt / \hbar)] = i\hbar \psi(\mathbf{r}) \left[-i \frac{E}{\hbar} \right] \exp(-iEt / \hbar) = E \Psi(\mathbf{r}, t) \end{aligned}$$

so $\Psi(\mathbf{r}, t) = \psi(\mathbf{r}) \exp(-iEt / \hbar)$ solves the
time-dependent Schrödinger equation

Compatibility with the time-independent equation

So every solution $\psi(\mathbf{r})$ of the time-independent Schrödinger equation, with eigenenergy E
is also a solution of the time-dependent equation
as long as we always multiply it by a factor $\exp(-iEt / \hbar)$

If $\psi(\mathbf{r})$ is a solution of the time-independent Schrödinger equation, with eigenenergy E

then $\Psi(\mathbf{r}, t) = \psi(\mathbf{r}) \exp(-iEt / \hbar)$

is a solution of both the time-independent
and the time-dependent Schrödinger equations
making these two equations compatible

Oscillations and time-independence

If we propose a solution

$$\Psi(\mathbf{r}, t) = \psi(\mathbf{r}) \exp(-iEt / \hbar)$$

to a time-independent problem


can this represent something that is stable in time?

Yes! - measurable quantities associated with this state

are stable in time!

e.g., probability density

$$|\Psi(\mathbf{r}, t)|^2 = \left[\exp(+iEt / \hbar) \psi^*(\mathbf{r}) \right] \times \left[\exp(-iEt / \hbar) \psi(\mathbf{r}) \right] = |\psi(\mathbf{r})|^2$$





8 The time-dependent Schrödinger equation

Slides: Lecture 8c Solutions of the time-dependent Schrödinger equation

Text reference: Quantum Mechanics for Scientists and Engineers

Section 3.3





The time-dependent Schrödinger equation



Solutions of the time-dependent Schrödinger equation

Quantum mechanics for scientists and engineers

David Miller

Contrast to classical wave equation

The common classical wave equation has a different form

$$\nabla^2 f = \frac{k^2}{\omega^2} \frac{\partial^2 f}{\partial t^2}$$

for which

$$f \propto \exp[i(kz - \omega t)]$$

would also be a solution

Note the classical equation has a second time derivative
as opposed to the first time derivative in
Schrödinger's time-dependent equation

Schrödinger's complex waves

Note that Schrödinger's use of a complex wave equation

$$-\frac{\hbar^2}{2m}\nabla^2\Psi(\mathbf{r},t)+V(\mathbf{r},t)\Psi(\mathbf{r},t)=i\hbar\frac{\partial\Psi(\mathbf{r},t)}{\partial t}$$

with the "i" on the right hand side

means that generally the wave Ψ is required to be a complex entity

For example, for $V = 0$

though $\exp\left[i(kz - Et/\hbar)\right]$ is a solution


$\sin(kz - Et/\hbar)$ is **not** a solution

Wave equation solutions

With the classical wave equation

if at some time we see a particular shape of wave

e.g., on a string



Wave equation solutions

With the classical wave equation

if at some time we see a particular shape of wave

e.g., on a string

we do not know if it is going

to the right $f(z - ct)$

Wave equation solutions

With the classical wave equation

if at some time we see a particular shape of wave

e.g., on a string

we do not know if it is going

to the right $f(z - ct)$

or to the left $g(z + ct)$

or even some combination of the two

Time evolution from Schrödinger's equation

In Schrödinger's equation, for a known potential V

$$-\frac{\hbar^2}{2m}\nabla^2\Psi(\mathbf{r},t)+V(\mathbf{r},t)\Psi(\mathbf{r},t)=i\hbar\frac{\partial\Psi(\mathbf{r},t)}{\partial t}$$

if we knew the wavefunction $\Psi(\mathbf{r},t_o)$ at every point in space at some time t_o

we could evaluate the left hand side of the equation at that time for all $\mathbf{r}$

so we would know $\partial\Psi(\mathbf{r},t)/\partial t$ for all $\mathbf{r}$

so we could integrate the equation to deduce $\Psi(\mathbf{r},t)$ at all future times

Time evolution from Schrödinger's equation

Explicitly

knowing $\partial\Psi(\mathbf{r},t)/\partial t$ we can calculate


$$\Psi(\mathbf{r},t_o + \delta t) \cong \Psi(\mathbf{r},t_o) + \left. \frac{\partial\Psi}{\partial t} \right|_{\mathbf{r},t_o} \delta t$$

that is, we can know the new wavefunction in space at the next instant in time

and we can continue on to the next instant

and so on

predicting all future evolution of the wavefunction





8 The time-dependent Schrödinger equation

Slides: Lecture 8d Linear superposition

Text reference: Quantum Mechanics for Scientists and Engineers

Sections 3.4 – 3.5





The time-dependent Schrödinger equation



Linear superposition

Quantum mechanics for scientists and engineers

David Miller

Linearity of Schrödinger's equation

The time-dependent Schrödinger equation is linear in the wavefunction Ψ

$$-\frac{\hbar^2}{2m}\nabla^2\Psi(\mathbf{r},t)+V(\mathbf{r},t)\Psi(\mathbf{r},t)=i\hbar\frac{\partial\Psi(\mathbf{r},t)}{\partial t}$$

One reason is that no higher powers of Ψ appear anywhere in the equation

A second reason is that Ψ appears in every term
there is no additive constant term anywhere

Linearity of Schrödinger's equation

Linearity requires two conditions

obeyed by Schrödinger's time-dependent equation

1 - If Ψ is a solution, then so also is $a\Psi$, where a is any constant

2 - If Ψ_a and Ψ_b are solutions, then so also is $\Psi_a + \Psi_b$

A consequence of these two conditions is that

$$\Psi_c(\mathbf{r}, t) = c_a \Psi_a(\mathbf{r}, t) + c_b \Psi_b(\mathbf{r}, t)$$

where c_a and c_b are (complex) constants

is also a solution

Linear superposition

The fact that

$$\Psi_c(\mathbf{r}, t) = c_a \Psi_a(\mathbf{r}, t) + c_b \Psi_b(\mathbf{r}, t)$$

is a solution if Ψ_a and Ψ_b are solutions

is the property of

linear superposition

To emphasize

linear superpositions of solutions of the
time-dependent Schrödinger equation
are also solutions

Time-dependence and expansion in eigenstates

We know that

if the potential V is constant in time

each of the energy eigenstates $\psi_n(\mathbf{r})$

with eigenenergy E_n

is separately a solution of the time-dependent Schrödinger equation

provided we remember to multiply by the right complex exponential factor

$$\Psi_n(\mathbf{r}, t) = \exp(-iE_n t / \hbar) \psi_n(\mathbf{r})$$

Time-dependence and expansion in eigenstates

Now we also know that the set of eigenfunctions of problems we will consider is a complete set

so the wavefunction at $t = 0$ can be expanded in them

$$\Psi(\mathbf{r}, 0) = \sum_n a_n \psi_n(\mathbf{r})$$

where the a_n are the expansion coefficients

But we know that a function that starts out as $\psi_n(\mathbf{r})$

will evolve in time as $\Psi_n(\mathbf{r}, t) = \exp(-iE_n t / \hbar) \psi_n(\mathbf{r})$

so, by linear superposition, the solution at time t is

$$\Psi(\mathbf{r}, t) = \sum_n a_n \Psi_n(\mathbf{r}, t) = \sum_n a_n \exp(-iE_n t / \hbar) \psi_n(\mathbf{r})$$

Time-dependence and expansion in eigenstates

Hence, for the case where the potential V does not vary in time

$$\Psi(\mathbf{r}, t) = \sum_n a_n \Psi_n(\mathbf{r}, t) = \sum_n a_n \exp(-iE_n t / \hbar) \psi_n(\mathbf{r})$$

is the solution of the time-dependent equation

with the initial condition $\Psi(\mathbf{r}, 0) = \psi(\mathbf{r}) = \sum_n a_n \psi_n(\mathbf{r})$

Hence, if we expand the wavefunction at time $t = 0$ in the energy eigenstates

we have solved for the time evolution of the state just by adding up the above sum

